

The Chvátal-Gomory Closure of a Strictly Convex Body is a Rational Polyhedron

Daniel Dadush and Santanu S. Dey
Georgia Institute of Technology

Juan Pablo Vielma
University of Pittsburgh

Inform's Annual Meeting. November, 2010 – Austin, TX

Outline


- Introduction
- Proof:
 - Step 1
 - Step 2
- Conclusions and Future Work

CG Cuts for Rational Polyhedra

$$P := \left\{ x \in \mathbb{R}^2 : \begin{array}{l} x_1 + x_2 \leq 3, \\ 5x_1 - 3x_2 \leq 3 \end{array} \right\}$$

$$\cap$$

$$H := \{ x \in \mathbb{R}^2 : 4x_1 + 3x_2 \leq 10.5 \}$$




CG Cuts for Rational Polyhedra

$$P := \left\{ x \in \mathbb{R}^2 : \begin{array}{l} x_1 + x_2 \leq 3, \\ 5x_1 - 3x_2 \leq 3 \end{array} \right\}$$

$$\cap$$

$$H := \{ x \in \mathbb{R}^2 : 4x_1 + 3x_2 \leq 10.5 \}$$




CG Cuts for Rational Polyhedra

$$P := \left\{ x \in \mathbb{R}^2 : \begin{array}{l} x_1 + x_2 \leq 3, \\ 5x_1 - 3x_2 \leq 3 \end{array} \right\}$$

$$\cap$$

$$H := \left\{ x \in \mathbb{R}^2 : \underbrace{4x_1 + 3x_2}_{\in \mathbb{Z}} \leq 10.5 \right\}$$

if $x \in \mathbb{Z}^2$



CG Cuts for Rational Polyhedra

$$P := \left\{ x \in \mathbb{R}^2 : \begin{array}{l} x_1 + x_2 \leq 3, \\ 5x_1 - 3x_2 \leq 3 \end{array} \right\}$$


$$\cap$$

$$H := \left\{ x \in \mathbb{R}^2 : \underbrace{4x_1 + 3x_2}_{\in \mathbb{Z}} \leq 10.5 \right\}$$

if $x \in \mathbb{Z}^2$

$$4x_1 + 3x_2 \leq [10.5]$$

Valid for $H \cap \mathbb{Z}^2$



CG Cuts for Rational Polyhedra

$$P := \left\{ x \in \mathbb{R}^2 : \begin{array}{l} x_1 + x_2 \leq 3, \\ 5x_1 - 3x_2 \leq 3 \end{array} \right\}$$

$$\cap$$


$$H := \left\{ x \in \mathbb{R}^2 : \underbrace{4x_1 + 3x_2}_{\in \mathbb{Z}} \leq 10.5 \right\}$$

if $x \in \mathbb{Z}^2$

$$4x_1 + 3x_2 \leq [10.5]$$

Valid for $H \cap \mathbb{Z}^2$

Valid for $C \cap \mathbb{Z}^2$



Importance of CG Cuts

- 3 main papers (Chvátal, Gomory, Schrijver)
 - +1,200 citations.
- First pure cutting plane algorithm for Integer Programming.
- CG cuts yield Matching Polytope.
- Cutting plane proofs.
- Still practical computationally.


CG Cuts for General Convex Sets

$$\sigma_C(a) := \sup\{\langle a, x \rangle : x \in C\}$$

$$C = \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$

$$\cap$$

$$H := \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$




CG Cuts for General Convex Sets

$$\sigma_C(a) := \sup\{\langle a, x \rangle : x \in C\}$$

$$C = \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$

$$\cap$$

$$H := \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$




CG Cuts for General Convex Sets

$$\sigma_C(a) := \sup\{\langle a, x \rangle : x \in C\}$$

$$C = \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$

$$H := \{x \in \mathbb{R}^n : \underbrace{\langle a, x \rangle}_{\in \mathbb{Z}} \leq \sigma_C(a)\}$$

if $x \in \mathbb{Z}^n$



CG Cuts for General Convex Sets

$$\sigma_C(a) := \sup\{\langle a, x \rangle : x \in C\}$$


$$C = \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$

$$H := \{x \in \mathbb{R}^n : \underbrace{\langle a, x \rangle}_{\in \mathbb{Z}} \leq \sigma_C(a)\}$$

if $x \in \mathbb{Z}^n$

$$\langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor$$

Valid for $H \cap \mathbb{Z}^n$



CG Cuts for General Convex Sets

$$\sigma_C(a) := \sup\{\langle a, x \rangle : x \in C\}$$

$$C = \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$


$$H := \{x \in \mathbb{R}^n : \underbrace{\langle a, x \rangle}_{\in \mathbb{Z}} \leq \sigma_C(a)\}$$

if $x \in \mathbb{Z}^n$

$$\langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor$$

Valid for $H \cap \mathbb{Z}^n$

Valid for $C \cap \mathbb{Z}^n$



CG Closure = Add all CG Cuts

$$\text{CGC}(C) := \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor\}$$

- Not necessarily a polyhedron, remember:

$$C = \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$

CG Closure = Add all CG Cuts

$$\text{CGC}(C) := \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor\}$$

- Not necessarily a polyhedron, remember:

$$C = \bigcap_{a \in \mathbb{Z}^n} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \sigma_C(a)\}$$

- $\text{CGC}(C)$ is a polyhedron if C is:
 - a rational polyhedra (Schrijver, 1980).
 - a “rational” ellipsoid (Dey and V. 2010).

Finite Number of Important CG Cuts

Theorem: There exists finite $S \subseteq \mathbb{Z}^n$ such that

$$\text{CGC}(C) = \bigcap_{a \in S} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor\}$$

Finite Number of Important CG Cuts

Theorem: There exists finite $S \subseteq \mathbb{Z}^n$ such that

$$\text{CGC}(C) = \bigcap_{a \in S} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor\}$$

- For this talk C is intersection of:
 - A strictly convex body (full dim, compact).
 - and a rational polyhedron.

Proof Outline for $\text{bd}(C) \cap \mathbb{Z}^n = \emptyset$

Theorem: There exists finite $S \subseteq \mathbb{Z}^n$ such that

$$\text{CGC}(C) = \text{CGC}(S, C) := \bigcap_{a \in S} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor\}$$

Proof Outline for $\text{bd}(C) \cap \mathbb{Z}^n = \emptyset$

Theorem: There exists finite $S \subseteq \mathbb{Z}^n$ such that

$$\text{CGC}(C) = \text{CGC}(S, C) := \bigcap_{a \in S} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor\}$$

- Step 1: There exists finite $S^1 \subseteq \mathbb{Z}^n$ such that
 - $\text{CGC}(S^1, C) \subseteq \text{int}(C)$

Proof Outline for $\text{bd}(C) \cap \mathbb{Z}^n = \emptyset$

Theorem: There exists finite $S \subseteq \mathbb{Z}^n$ such that


$$\text{CGC}(C) = \text{CGC}(S, C) := \bigcap_{a \in S} \{x \in \mathbb{R}^n : \langle a, x \rangle \leq \lfloor \sigma_C(a) \rfloor\}$$

- Step 1: There exists finite $S^1 \subseteq \mathbb{Z}^n$ such that
 - $\text{CGC}(S^1, C) \subseteq \text{int}(C)$
- Step 2: There exists finite $S^2 \subseteq \mathbb{Z}^n$ such that
 - $\text{CGC}(C) = \text{CGC}(S^1, C) \cap \text{CGC}(S^2, C)$

Separate points in $\text{bd}(C)$

$$u \in \text{bd}(C) \setminus \mathbb{Z}^n \quad \exists a^u \in \mathbb{Z}^n$$

$$\langle a^u, u \rangle > \lfloor \sigma_C(a^u) \rfloor$$




Separate points in $\text{bd}(C)$

$$u \in \text{bd}(C) \setminus \mathbb{Z}^n \quad \exists a^u \in \mathbb{Z}^n$$

$$\langle a^u, u \rangle > \lfloor \sigma_C(a^u) \rfloor$$

$$\langle s(u), u \rangle = \sigma_C(s(u))$$




Separate points in $\text{bd}(C)$

$$u \in \text{bd}(C) \setminus \mathbb{Z}^n \quad \exists a^u \in \mathbb{Z}^n$$

$$\langle a^u, u \rangle > \lfloor \sigma_C(a^u) \rfloor$$

$$\underbrace{\langle s(u), u \rangle}_{\in \mathbb{Z}^n} = \underbrace{\sigma_C(s(u))}_{\notin \mathbb{Z}}$$




Separate points in $\text{bd}(C)$

$$u \in \text{bd}(C) \setminus \mathbb{Z}^n \quad \exists a^u \in \mathbb{Z}^n$$

$$\langle a^u, u \rangle > \lfloor \sigma_C(a^u) \rfloor$$

$$\underbrace{\langle s(u), u \rangle}_{\in \mathbb{Z}^n} = \underbrace{\sigma_C(s(u))}_{\in \mathbb{Z}}$$



Separate points in $\text{bd}(C)$

$$u \in \text{bd}(C) \setminus \mathbb{Z}^n \quad \exists a^u \in \mathbb{Z}^n$$

$$\langle a^u, u \rangle > \lfloor \sigma_C(a^u) \rfloor$$

$$\underbrace{\langle s(u), u \rangle}_{\in \mathbb{Z}^n} = \underbrace{\sigma_C(s(u))}_{\in \mathbb{Z}} \quad \checkmark$$

$$\lambda s(u) \notin \mathbb{Z}^n \quad \forall \lambda > 0 :$$

$$\lambda s(u) \in \mathbb{Z}^n \Rightarrow \sigma_C(\lambda s(u)) \in \mathbb{Z} :$$

Separate points in $\text{bd}(C)$

$$u \in \text{bd}(C) \setminus \mathbb{Z}^n \quad \exists a^u \in \mathbb{Z}^n$$

$$\langle a^u, u \rangle > \lfloor \sigma_C(a^u) \rfloor$$

$$\underbrace{\langle s(u), u \rangle}_{\in \mathbb{Z}^n} = \underbrace{\sigma_C(s(u))}_{\notin \mathbb{Z}}$$



$$\lambda s(u) \notin \mathbb{Z}^n \quad \forall \lambda > 0 :$$

$$C = \left\{ x \in \mathbb{R}^2 : \sqrt{x_1^2 + x_2^2} \leq 1 \right\}$$

$$u = (1/2, \sqrt{3}/2)^T \in \text{bd}(C)$$

$$s(u) = (1/2, \sqrt{3}/2)^T$$

$$\lambda s(u) \in \mathbb{Z}^n \Rightarrow \sigma_C(\lambda s(u)) \in \mathbb{Z} :$$

Separate points in $\text{bd}(C)$

$$u \in \text{bd}(C) \setminus \mathbb{Z}^n \quad \exists a^u \in \mathbb{Z}^n$$

$$\langle a^u, u \rangle > \lfloor \sigma_C(a^u) \rfloor$$

$$\underbrace{\langle s(u), u \rangle}_{\in \mathbb{Z}^n} = \underbrace{\sigma_C(s(u))}_{\notin \mathbb{Z}}$$



$$\lambda s(u) \notin \mathbb{Z}^n \quad \forall \lambda > 0 :$$

$$C = \left\{ x \in \mathbb{R}^2 : \sqrt{x_1^2 + x_2^2} \leq 1 \right\}$$

$$u = (1/2, \sqrt{3}/2)^T \in \text{bd}(C)$$

$$s(u) = (1/2, \sqrt{3}/2)^T$$

$$\lambda s(u) \in \mathbb{Z}^n \Rightarrow \sigma_C(\lambda s(u)) \in \mathbb{Z} :$$

$$C = \left\{ x \in \mathbb{R}^2 : \sqrt{x_1^2 + x_2^2} \leq 5 \right\}$$

$$u = (25/13, 60/13)^T \in \text{bd}(C)$$

$$s(u) = (5, 12)^T, \quad \sigma_C(s(u)) = 65$$

Separate points in $\text{bd}(C)$

$$u \in \text{bd}(C) \setminus \mathbb{Z}^n \quad \exists a^u \in \mathbb{Z}^n$$

$$\langle a^u, u \rangle > \lfloor \sigma_C(a^u) \rfloor$$

$$\underbrace{\langle s(u), u \rangle}_{\in \mathbb{Z}^n} = \underbrace{\sigma_C(s(u))}_{\notin \mathbb{Z}}$$



$$\frac{s^i}{\|s^i\|} \xrightarrow{i \rightarrow \infty} \frac{s(u)}{\|s(u)\|}$$

$$\lim_{i \rightarrow \infty} \langle s^i, u \rangle - \lfloor \sigma_C(s^i) \rfloor > 0$$

Diophantine approx. of $s(u)$

$$\lambda s(u) \notin \mathbb{Z}^n \quad \forall \lambda > 0 :$$

$$C = \left\{ x \in \mathbb{R}^2 : \sqrt{x_1^2 + x_2^2} \leq 1 \right\}$$

$$u = (1/2, \sqrt{3}/2)^T \in \text{bd}(C)$$

$$s(u) = (1/2, \sqrt{3}/2)^T$$

$$\lambda s(u) \in \mathbb{Z}^n \Rightarrow \sigma_C(\lambda s(u)) \in \mathbb{Z} :$$

$$C = \left\{ x \in \mathbb{R}^2 : \sqrt{x_1^2 + x_2^2} \leq 5 \right\}$$


$$u = (25/13, 60/13)^T \in \text{bd}(C)$$

$$s(u) = (5, 12)^T, \sigma_C(s(u)) = 65$$

Compactness Argument

$$K := \text{bd}(C)$$

$$S_u := \{x : \langle a^u, x \rangle > \lfloor \sigma_C(a^u) \rfloor\}$$




Compactness Argument

$$K := \text{bd}(C)$$

$$S_u := \{x : \langle a^u, x \rangle > \lfloor \sigma_C(a^u) \rfloor\}$$

$$K \subset \bigcup_{u \in K} S_u$$




Compactness Argument

$$K := \text{bd}(C)$$

$$S_u := \{x : \langle a^u, x \rangle > \lfloor \sigma_C(a^u) \rfloor\}$$

$$\text{compact} \longrightarrow K \subset \bigcup_{u \in K} S_u$$



Compactness Argument


$$K := \text{bd}(C)$$

$$S_u := \{x : \langle a^u, x \rangle > \lfloor \sigma_C(a^u) \rfloor\}$$

$$\text{compact} \longrightarrow K \subset \bigcup_{u \in K} S_u$$

$$\downarrow$$

$$K \subset \bigcup_{i=1}^m S_{u^i}$$



Compactness Argument

$$K := \text{bd}(C)$$


$$S_u := \{x : \langle a^u, x \rangle > \lfloor \sigma_C(a^u) \rfloor\}$$

compact \longrightarrow $K \subset \bigcup_{u \in K} S_u$

\downarrow

$$K \subset \bigcup_{i=1}^m S_{u^i}$$


$$S^1 = \bigcup_{i=1}^m \{a^{u^i}\}$$



Step 2 : Separate $\text{CGC}(S^1, C) \setminus \text{CGC}(C)$

$$V := \text{Ext}(\text{CGC}(S^1, C)) \setminus \mathbb{Z}^n$$

$$\langle a, v \rangle > \lfloor \sigma_C(a) \rfloor$$




Step 2 : Separate $\text{CGC}(S^1, C) \setminus \text{CGC}(C)$

$$V := \text{Ext}(\text{CGC}(S^1, C)) \setminus \mathbb{Z}^n$$

$$\langle a, v \rangle > \lfloor \sigma_C(a) \rfloor$$

$$\exists \varepsilon > 0 \quad \varepsilon B^n + v \subset C \quad \forall v \in V$$



Step 2 : Separate $\text{CGC}(S^1, C) \setminus \text{CGC}(C)$

$$V := \text{Ext}(\text{CGC}(S^1, C)) \setminus \mathbb{Z}^n$$

$$\langle a, v \rangle > \lfloor \sigma_C(a) \rfloor$$

$$\exists \varepsilon > 0 \quad \varepsilon B^n + v \subset C \quad \forall v \in V$$


$$\|a\| \geq \frac{1}{\varepsilon} \Rightarrow$$

$$\lfloor \sigma_C(a) \rfloor \geq \sigma_C(a) - 1$$

$$\geq \sigma_{v+\varepsilon B^n}(a) - 1$$

$$= \langle v, a \rangle + \varepsilon \|a\| - 1$$

$$\geq \langle v, a \rangle$$



Step 2 : Separate $\text{CGC}(S^1, C) \setminus \text{CGC}(C)$

$$V := \text{Ext}(\text{CGC}(S^1, C)) \setminus \mathbb{Z}^n$$


$$\langle a, v \rangle > \lfloor \sigma_C(a) \rfloor$$

$$\exists \varepsilon > 0 \quad \varepsilon B^n + v \subset C \quad \forall v \in V$$

$$\|a\| \geq \frac{1}{\varepsilon} \Rightarrow$$

$$\begin{aligned} \lfloor \sigma_C(a) \rfloor &\geq \sigma_C(a) - 1 \\ &\geq \sigma_{v+\varepsilon B^n}(a) - 1 \\ &= \langle v, a \rangle + \varepsilon \|a\| - 1 \\ &\geq \langle v, a \rangle \end{aligned}$$

$$S^2 = (1/\varepsilon)B \cap \mathbb{Z}^n$$



Conclusions and Future Work

- Non-Constructive because of compactness argument in step 1.
- General compact convex sets including non-rational polytopes done:
 - Dadush, Dey and V. MB39: Monday 11:00, C - Room 9B, Level 3
- Open Problems:
 - Simpler Proof (Circle in \mathbb{R}^2 ?).
 - Constructive/Algorithmic proof.